



HERBACROWN

720g/L 2-4-D amine salt



PHI: 10-12 days

Systemic selective herbicide (growth inhibitor) for the control of broadleaved weeds in maize, millet, rice, sugarcane, sorghum. It can also be used to kill trees.

DIRECTIONS FOR USE

- Avoiding overdosing 2,4-D during weed control in crops, to ensure crops are not damaged from its growth inhibiting function.
- Cereals (Wheat, Oats, Rice, Barley): Treatment with 2,4-D Amine salt should occur at some time between tillering and the end of shooting. 2,4-D should not be used at any later stage, as it will interfere with fruiting. Oats are more sensitive to 2,4-D than the other cereal types.
- Rice: Rice can be treated with 2,4-D Amine when the plants are fully established, 2-3 weeks after emergence, depending on the variety and the climatic conditions. Transplanted rice can be treated when the plants are fully established. The field/flooded water should be drained 24 hours before application. The field may be re-flooded three days after application to enhance the dying process.
- Millet (sorghum): Millet can be treated with 2,4-D Amine when it has grown to a height of between 10 and 25cm.
- Maize: In maize field, application of 2,4-D can be carried out at two times: before emergence or when the maize is between 10 and 25cm tall. Due to the "shock" effect from 2,4-D, spraying of maize above 25cm height will cause lodging of the maize plant and affect plant development. Spraying 2,4-D in maize at tasseling and fruiting stage will inhibit growth and interfere with fruiting.

APPLICATION GUIDE

- Ensure to use a well functioning sprayer
- Use clean water, free of debris (sieve appropriately)
- Fix and use 2.4m diameter nozzle (Flood).
- Double-rinse the knapsack sprayer and lance by pumping out clean water.
- Half fill the tank with water and measure the appropriate dosage into the knapsack.
- For tank-mixed application with other active ingredient(s), add the appropriate dosage of the other product. *Add sticker if rainfall is expected in less than 6 hours.*
- Fill the sprayer tank with water to 15L and mix the solution by pumping and spraying into the tank, while stirring.
- Ensure there is no leakage before mounting the knapsack sprayer.
- Pump out spray solution whilst maintaining normal pressure for a fine spray mist.

DOSAGE:

50ml-80ml (weed control in crops), 100ml-150ml (weed control in non-cropped areas) per 15L knapsack



MANUFACTURED BY

Agro Fortune Company Limited

ORIGIN: CHINA

DISTRIBUTED BY

Agrocrown West Africa Co. Ltd.

First Floor, Ivory Building, 22 Nii

Martey Tsuru St, Spintex Road,

Accra

info@agrocrown.com



www.agrocrown.com



+233 34 229 5994



AgroCrown
Farmers' Choice